

# Radial Velocity Variations in Pulsating Ap Stars.

## III. The Discovery of 16.21 min Oscillations in $\beta$ CrB $\star$

A.P. Hatzes<sup>1</sup> and D.E. Mkrtichian<sup>2,3</sup>

<sup>1</sup> *Thüringer Landessternwarte Tautenburg, Sternwarte 5, D-07778, Tautenburg, Germany*

<sup>2</sup> *Astrophysical Research Center for the Structure and Evolution of the Cosmos, Sejong University, Seoul 143-747, Korea*

<sup>3</sup> *Astronomical Observatory, Odessa National University, Shevchenko Park, Odessa, 65014, Ukraine*

### ABSTRACT

We present the analysis of 3 hrs of a rapid time series of precise stellar radial velocity (RV) measurements ( $\sigma = 4.5 \text{ m s}^{-1}$ ) of the cool Ap star  $\beta$  CrB. The integrated RV measurements spanning the wavelength interval 5000–6000 Å show significant variations (false alarm probability =  $10^{-5}$ ) with a period of 16.21 min ( $\nu = 1028.17 \mu\text{Hz}$ ) and an amplitude of  $3.54 \pm 0.56 \text{ m s}^{-1}$ . The RV measured over a much narrower wavelength interval reveals one spectral feature at  $\lambda 6272.0 \text{ Å}$  pulsating with the same 16.21 min period and an amplitude of  $138 \pm 23 \text{ m s}^{-1}$ . These observations establish  $\beta$  CrB to be a low-amplitude rapidly oscillating Ap star.

**Key words:** Stars:individual; $\beta$  CrB – Stars:pulsation – Stars:variables

## 1 INTRODUCTION

Kurtz (1989) first suggested that the cool magnetic Ap star  $\beta$  CrB was a prime candidate to be a rapidly oscillating Ap (roAp) star since it had stellar properties similar to several known oscillating Ap stars (HR 1217, 33 Lib, and  $\gamma$  Equ). The issue of whether  $\beta$  CrB is a pulsating star has important implications for the excitation mechanism in roAp stars as well as the possible existence of an instability strip for roAp stars.

Several investigators have searched for pulsational variations in  $\beta$  CrB using photometric measurements with null results (Heller & Kramer 1988; Kreidl 1991). Recently, Kochukhov et al. (2002, hereafter K02) presented radial velocity (RV) measurements for  $\beta$  CrB spanning the wavelength interval 6105 – 6190 Å. Measurements of most lines were constant to a level of  $20\text{--}30 \text{ m s}^{-1}$ ; however, intriguing evidence for RV variations were found for one spectral feature, Fe I  $\lambda 6165.4 \text{ Å}$ . This feature had an amplitude of  $71 \pm 11 \text{ m s}^{-1}$  and a period of  $11.5 \pm 0.5 \text{ min}$ . Before we can place  $\beta$  CrB in the family of roAp stars the K02 result must be confirmed since the false alarm probability for the detection was rather high ( $= 0.016$ ). Here we present our own precise radial velocity measurements for  $\beta$  CrB and show that it is indeed a rapidly oscillating Ap star, but not with the period found by K02.

## 2 OBSERVATIONS

Spectral observations of  $\beta$  CrB were made for 3 consecutive hours on the night of 31 January 1998 (JD = 2450844.903220) using the “2dcoude” echelle spectrograph of the Harlan J. Smith 2.7m telescope of McDonald Observatory (Tull et al. 1995). According to the ephemeris of Kurtz (1989) this corresponds to magnetic phase 0.11. One hundred fifty-five observations were made with exposure times of 50 secs which yielded a typical signal-to-noise ratio of 150 per pixel. The description of the instrumental setup can be found in previous papers (Kanaan & Hatzes 1998; Mkrtichian et al. 2003). Briefly, the wavelength coverage was 4700 – 6700 Å at a resolving power of 60,000. Precise stellar radial velocities were achieved via an iodine absorption cell placed in the optical light path.

## 3 RESULTS

### 3.1 Integrated RV Measurements

To achieve the maximum RV measurement precision we first used the full spectral region covered by the iodine reference absorption lines to produce a “mean” RV value for all spectral lines. In doing so we excluded the two reddest spectral orders that covered the wavelength range 6100–6300 Å. Here the iodine absorption lines are weak and this degrades the RV precision. To determine the relative RV for  $\beta$  CrB each spectral order was divided into 60 chunks so that we could model the spatial (and temporal) variations of the instrumental profile (IP) which can introduce significant RV er-




**Figure 1.** (Top) The Lomb-Scargle (L-S) periodogram of the RV measurements of  $\beta$  CrB in the wavelength interval 5000–6000 Å. The L-S power (unitless) is a measure of statistical significance. (Bottom) The discrete Fourier transform (amplitude spectrum) of the same RV measurements. In this case amplitude is in units of  $\text{m s}^{-1}$ .

rors. In each chunk the RV shift was calculated along with the IP using the IP reconstruction procedure outlined in Valenti et al. (1996). The RV measurements from all chunks were then combined weighted by the inverse square of the RV standard deviation for each chunk. RV measurements of an individual chunk that differed by more than  $4\sigma$  from the mean were rejected in the calculation of the mean.

The RV measurements had a long-term trend of unknown origin. This could be due to rotational modulation caused by the abundance spots known to occur on Ap stars, long-period oscillations, or a stellar or sub-stellar companion. Observations with a longer timebase are needed to answer this. Since we are primarily interested in searching for pulsational variations on time scales of roAp stars (5–15 min), a second order polynomial was fitted to the long-term variations and subtracted. Variations in the sky transparency during the course of the observations resulted in some spectra having significantly lower signal-to-noise ratio ( $S/N$ ) than the mean ( $S/N \approx 150$ ). Observations having  $S/N < 100$  (fifteen data points) were eliminated from the time series analysis.

Figure 1 shows both the Lomb-Scargle (L-S) periodogram (Lomb 1976; Scargle 1982) and the discrete Fourier transform (DFT) of the residual RV time series for  $\beta$  CrB. These show significant power at  $1028.17 \mu\text{Hz}$  ( $P = 16.21$  min). When trying to detect a periodic signal in time series data *and* assessing its statistical significance the Lomb-Scargle (L-S) periodogram is a more appropriate analysis tool over the DFT. In this case the L-S power is not a mea-

sure of the amplitude of the signal, but instead is related to the statistical significance of a peak. Many researchers in the field of oscillating stars are more accustomed to dealing with DFT. In this instance the power is directly related to the true amplitude of the periodic signal in the data and can also be used to assess the noise level of the data. The DFT, however, has limited use in quantifying the significance of a signal. We thus show both the L-S periodogram and DFT (lower panel of Figure 1).

The L-S periodogram of the  $\beta$  CrB data has power more than 15 which is the first hint that we are dealing with a statistically significant signal. Using the Eq. 14 of Scargle (1982) this power results in a false alarm probability (FAP), i.e. the chance that noise is creating the signal, of  $\approx 2 \times 10^{-5}$ .

The power in a L-S periodogram only gives a rough estimate of the false alarm probability of a signal. A more accurate determination is to use a bootstrap randomization technique (e.g. Kürster et al. 1997). This technique randomly shuffles the RV measurements keeping the times fixed. A Lomb-Scargle periodogram is calculated for each “random” data set and these data re-shuffled. After a large number of these shuffles, the fraction of the periodograms having maximum power greater than the observed data periodogram over the frequency interval of interest (in this case  $\nu = 200\text{--}5000 \mu\text{Hz}$ ) represents the false alarm probability (FAP), or the chance that random noise can produce the observed power in the periodogram. Using  $2 \times 10^5$  shuffles we establish a  $\text{FAP} = 10^{-5}$  for the peak in Figure 1.

Figure 2 shows the residual RV measurements after removal of the long-term trend phased to the 16.2 min period. The top panel is the raw data while the bottom panel represents phase-binned ( $\Delta\phi \approx 0.05$ ) values. The amplitude of these RV variations is  $3.54 \pm 0.56 \text{ m s}^{-1}$  ( $K$ -amplitude).

### 3.2 Narrow-band Measurements

As first shown by Kanaan & Hatzes (1998) the RV amplitude of individual spectral lines in roAp stars can differ by factors of 10–100. Later studies established that elements of Nd and Pr in some roAp stars had the highest pulsational RV amplitudes (Savanov et al. 1999, Kochukhov & Ryabchikova 2001; Kurtz et al. 2003). To search for spectral lines that that might have high amplitude RV variations the individual wavelength chunks (each chunk typically contained at most 1–2 spectral lines) prior to combining these in a mean velocity were examined. Because high RV amplitude lines can also be found in the reddest orders (excluded from the integrated RV analysis) these were also used to search for RV variations). A periodogram analysis was performed on the RVs from the individual wavelength chunks. If a periodogram showed significant power (Lomb-Scargle power  $> 7$ ) a statistical assessment was made using the bootstrap randomization technique (50,000–200,000 shuffles).

Only one chunk ( $\lambda 6271.7 - \lambda 6273.3 \text{ Å}$ ) had highly significant variations with the same period as found in the integrated RV data. The spectral region for this chunk is shown in Figure 3. The RV amplitude ( $K$ ) for this chunk,  $138 \pm 23 \text{ m s}^{-1}$ , was significantly higher than the value for the integrated RV measurements. Figure 4 shows both the Lomb-Scargle periodogram and DFT of the RVs from this wavelength chunk. Using 200,000 shuffles we established  $\text{FAP} \approx 1.5 \times 10^{-5}$  for this peak. The pulsational phase for this




**Figure 2.** (Top) The residual RV variations phased to a period of 16.2 min. (Bottom) The phased-binned averages ( $\Delta\phi \approx 0.05$ ) for data shown in the top panel. In both panels data points are repeated for the second cycle.




**Figure 3.** Spectral region of the wavelength chunk ( $\lambda 6271.7 - \lambda 6273.3$  Å) showing high amplitude RV variations.




**Figure 4.** The Lomb-Scargle periodogram (top) and DFT (bottom) of the RV variations for the spectral region ( $\lambda 6271.7 - \lambda 6273.3$  Å)

chunk is identical to within the errors of the pulsation phase calculated using the integrated RV measurements.

We tentatively identify the spectral features in the high amplitude chunk as being a Ce II  $\lambda 6272.026$  Å line that is blended in the blue wing by a weaker Cr II  $\lambda 6271.870$  Å feature.

## 4 DISCUSSION

Our precise RV measurements for  $\beta$  CrB establish that this star is with high probability a rapidly oscillating Ap star with an amplitude of  $3.54 \pm 0.56$   $\text{m s}^{-1}$  and a period of  $16.2 \pm 0.7$  min. Although additional observations are needed to confirm our result, we believe that the signal we have detected is real for a number of reasons:

- *We derive a very low false alarm probability.*

The probability that this is a false signal due to noise is  $\approx 10^{-5}$  as determined through the bootstrap randomization procedure. This is a more rigorous way of determining the FAP than strictly using the Lomb-Scargle power. This result represents the combined mean velocity of all spectral lines spanning the wavelength interval 5000–6000 Å. Any systematic or instrumental error would have to affect all wavelength chunks used in the analysis in the same way which seems unlikely.

- *The variations are not due to a few outliers.*

We think it is unlikely that a few outliers are causing our signal. (We define an outlier as deviating significantly from the mean RV value and not from the best fit sine wave through the data.) If only a few outliers are driving the

power in the periodogram then this would be evident in a high false alarm probability.

For a real periodic signal in the presence of noise, the more data one accumulates the more significant the detection becomes, even with the presence of a few occasional outliers. Thus the power in a Lomb-Scargle periodogram should increase with increasing number of data points. This is demonstrated in Figure 5 which shows the Lomb-Scargle power at  $\nu = 1028.17 \mu\text{Hz}$  (circles) as a function of the number of data points used in the periodogram. For comparison we generated a synthetic signal consisting of a sine wave with the same period and amplitude as found in our data and sampled in the same manner as the real data. Random noise was added with the same rms scatter as our measurements. The crosses in Figure 5 show that the L-S power of the fake data (with a signal present) increases with increasing number of data points in the same way.

An iodine absorption cell was used to perform our RV measurements because it is designed to eliminate, or at least minimize instrumental effects. However, one could argue that there might be residual instrumental shifts not corrected by the reduction process, or that our instrumental profile modelling somehow introduces a false signal into the data. We do not believe this is the case.

We investigated whether periodic instrumental shifts were present in our data using the wavelength solution calculated from the iodine absorption lines. A Fourier analysis of these shifts showed the strongest period at 176 minutes, and smaller variations with periods of 20 and 24 minutes. All periods were significantly different from the one found in our RV data. Furthermore, we also performed a Fourier analysis of all IP parameters used in our RV determination. We found no variations at the period coinciding with the one found in the RV data. This demonstrates that using the iodine cell technique does an excellent job of excluding instrumental variations.

- *The same signal is found in the  $\lambda 6272 \text{ \AA}$  feature.*

The spectral orders used to determine the integrated RV did not include the order covering the  $\lambda 6272 \text{ \AA}$  feature. Thus this spectral line is not contributing to the signal found in our integrated RV measurements. However, a separate analysis of this spectral chunk did find the same 16.21 min period and with the same statistical significance as with the integrated RV measurements. Although this is not an independent confirmation (we are using the same data set, but different parts of the spectrum), this result and the above arguments argue strongly that the 16.21 min oscillations in  $\beta \text{ CrB}$  are real.

Our RV measurements fail to confirm the results of K02 on the Fe I  $\lambda 6165 \text{ \AA}$  feature. The wavelength chunk containing this spectral line had an rms scatter of  $130 \text{ m s}^{-1}$  (due to the weak iodine absorption lines in this wavelength region). However, we had a factor of 3 more measurements than K02. Monte Carlo simulations indicate that we would have detected an 11.5 min period with an amplitude of  $70 \text{ m s}^{-1}$  and a FAP of  $10^{-4}$ . An amplitude of  $50 \text{ m s}^{-1}$  would have been detected with a FAP = 0.01. Thus we would have detected any variations in the Fe I  $\lambda 6165 \text{ \AA}$  line if they were present at the same amplitude as reported by K02.

We believe that the result of K02 on  $\beta \text{ CrB}$  is spurious. The FAP is much too high (FAP = 0.016) and our measure-




**Figure 5.** The Lomb-Scargle power at  $\nu = 1028.17 \mu\text{Hz}$  as a function of number of data points used in the periodogram using real (circles) and fake data (crosses). The fake data was generated using a sine wave with the same amplitude ( $3.56 \text{ m s}^{-1}$ ) and period ( $= 16.21 \text{ min}$ ) as found in our data and sampled in the same manner. Random noise with the same scatter ( $\sigma = 4.5 \text{ m s}^{-1}$ ) as our data was also added to the fake signal.

ments have failed to find the presence of an 11.5 min period in either the integrated RV measurements, or in the analysis of the narrow wavelength chunks. More troubling is that the K02 period of  $11.5 \pm 0.5 \text{ min}$  found in  $\beta \text{ CrB}$  is very close to the values of periods ( $\approx 11.7 \text{ min}$ ) found in several spectral lines in the roAp star 10 Aql using the same instrument.

The most likely explanation for the RV signal found in  $\beta \text{ CrB}$  by K02 is that it is due to short term uncorrected instrumental variations in the Gecko spectrograph used for the measurements. Circumstantial evidence for this comes from Matthews & Scott (1995) who reported an 11.1 min period in the RV variations in  $\gamma \text{ Equ}$ . This period is different from known pulsation modes and has never been confirmed by subsequent RV measurements from other investigations. Although the authors used a superimposed mercury emission line from an arc lamp to eliminate instrumental shifts, the 11.1 min period could still be a residual instrumental effect due to the stellar light a calibration source having a slightly different optical path (not the case when using the iodine absorption cell).

The RV measurements of K02, on the other hand, were made *without* a simultaneous wavelength calibration (either  $\text{I}_2$  absorption cell or simultaneous Th-Ar calibration), rather the wavelength calibration used a Th-Ar exposure taken before and after the time series used for the RV measurements. This cannot correct for any instrumental variations on time scales significantly shorter than the time between the two calibration measurements. Because K02 have failed to ex-

clude an instrumental origin with a period of  $\approx 11.5$  min for the RV variations in  $\beta$  CrB their results on this star (and possibly 10 Aql) should be considered suspect. An investigation of the short-term instrumental shifts of the Gecko spectrograph would be useful.

Our RV measurements for  $\beta$  CrB seem to establish that this star is an roAp star with the lowest RV amplitude ( $3.5 \text{ m s}^{-1}$ ) and one of the longest periods (16.2 min).  $\beta$  CrB is also unique among roAp stars in that it shows high amplitude pulsational in only one spectral feature, the  $\lambda 6271.9 \text{ \AA}$ . This showed significant variations ( $\text{FAP} = 1.5 \times 10^{-5}$ ) with the same 16.2 min period and an amplitude of  $138 \pm 23 \text{ m s}^{-1}$ . We tentatively identify this feature as a blend of Ce II and Cr II lines. Why  $\beta$  CrB has such a low RV amplitude, almost no high amplitude spectral lines, and a longer period mode compared to other roAp stars with similar spectral properties should provide clues as to the origin of the roAp phenomenon. We are currently planning observations of this star over a full rotation period not only to confirm our detection, but to search for rotationally modulated amplitude variations of the pulsations.

## 5 ACKNOWLEDGEMENTS

APH acknowledges the support of grant 50OW0204 from the Deutsches Zentrum für Luft- und Raumfahrt e.V. (DLR). DEM acknowledges his work as part of research activity of the Astrophysical Research Center for the Structure and Evolution of the Cosmos (ARCSEC) which is supported by the Korean Science and Engineering Foundation. We thank William Cochran for useful discussions and for his assistance in developing the radial velocity reduction software and Antonio Kanaan for his help in acquiring the data.

## REFERENCES

- Heller, C.H., & Kramer, K.S. 1988, *PASP*, 238, 261.  
 Kanaan, A., Hatzes, A.P. 1998, *ApJ*, 503, 848  
 Kreidl, T.J. 1991, *MNRAS* 248, 701  
 Kochukhov, O., Ryabchikova, T. 2001, *A&A*, 374, 615.  
 Kochukhov, O., Landstreet, J. D., Ryabchikova, T., Weiss, W. W. & Kupka, F. 2002, *MNRAS*, 337, L1 (K02)  
 Kürster, M., Schmitt, J.H.M.M., Cutispoto, G. & Dennerl, K. 1997 *A&A* 320, 831  
 Kurtz, D.W. 1989, *MNRAS* 238, 261  
 Kurtz, D.W., Elkin, V.G., & Mathys, G. 2003, 343, L5.  
 Lomb N.R. 1976, *Ap&SS*, 39, 477  
 Matthews J.,M. Scott, S. 1995, in *Astrophysical Applications of Stellar Oscillations*, eds. R.S. Stobie and P.A. Whitelock, ASP Conf. Series, 83, p347  
 Mkrtichian, D.E., Hatzes A.P., Kanaan A., 2003, *MNRAS*, 345, 781  
 Savanov, I.S., Malanushenko, V.P., & Ryabchikova, T.A. 1999, *Astronomy Letters*, 25, 902.  
 Scargle, J. D. 1982, *ApJ*, 263, 835  
 Tull, R.G., MacQueen, P.J., Sneden, C. & Lambert, D.L. 1995, *PASP*, 107, 251.  
 Valenti, J. A., Butler, R. P., & Marcy, G. W. 1995, *PASP*, 107, 966